

AD

RSIC-647

**REPORT 4: INSTABILITY OF PERIODIC
MOTIONS NEAR COLLINEAR POINTS OF LIBRATION
IN THE RESTRICTED PROBLEM OF THREE BODIES**

by

G. N. Duboshin

Astronomicheskii Zhurnal, 15, No. 3, 209-216 (1938)

Translated from the Russian

February 1967

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

REDSTONE SCIENTIFIC INFORMATION CENTER
REDSTONE ARSENAL, ALABAMA

JOINTLY SUPPORTED BY



U.S. ARMY MISSILE COMMAND



GEORGE C. MARSHALL SPACE FLIGHT CENTER

N67-26282
(ACCESSION NUMBER)
12
(PAGES)
JMX-59661
(NASA CR OR TMX OR AD NUMBER)

(THRU)
(CODE)
30
(CATEGORY)

Disclaimer

The findings of this report are not to be construed as an official Department of the Army position.

Disposition

Destroy this report when it is no longer needed. Do not return it to the originator.

13 February 1967

RSIC-647

**REPORT 4: INSTABILITY OF PERIODIC
MOTIONS NEAR COLLINEAR POINTS OF LIBRATION
IN THE RESTRICTED PROBLEM OF THREE BODIES**

by

G. N. Duboshin

Astronomicheskii Zhurnal, 15, No. 3, 209-216 (1938)

(The First Three Reports from the Group of Papers
Entitled "The Gil'den-Mul'ton Hypothesis on the
Origin of Counter-Glow" Were Published in the
Preceding Issue of This Journal)

Translated from the Russian

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Translation Branch
Redstone Scientific Information Center
Research and Development Directorate
U. S. Army Missile Command
Redstone Arsenal, Alabama 35809

This article examines the problem of periodic motions near collinear points of libration in the plane restricted problem of three bodies. With the aid of the Lyapunov methods, we demonstrate that around each collinear point of libration, a continuous family of periodic motions exists, dependent on one arbitrary constant. We examine the stability of the indicated periodic motions and show that they are unstable. Finally, we establish the orbital instability of the periodic motions under study.

We shall proceed from differential equations, determining the movement of a particle (zero mass) near any collinear point of libration. To be definite, we shall examine point L_2 . The indicated equations can be written as follows:

$$\left. \begin{aligned} \frac{d^2 \xi}{dt^2} - 2n \frac{d\eta}{dt} &= \frac{\partial \Omega}{\partial \xi}, \\ \frac{d^2 \eta}{dt^2} + 2n \frac{d\xi}{dt} &= \frac{\partial \Omega}{\partial \eta}, \end{aligned} \right\} \quad (1)$$

where ξ and η -- particle coordinates in a system of coordinates with its beginning at point L_2 , $n^2 = 1 + \mu$ and Ω is determined by formula

$$2\Omega = \rho_1^2 + \frac{2}{\rho_1} + \mu \left(\rho_2^2 + \frac{2}{\rho_2} \right), \quad (2)$$

where

$$\left. \begin{aligned} \rho_1^2 &= \left(\xi + a - \frac{\mu}{1+\mu} \right)^2 + \eta^2, \\ \rho_2^2 &= \left(\xi + a + \frac{1}{1+\mu} \right)^2 + \eta^2. \end{aligned} \right\} \quad (3)$$

Finally, $|a|$ is the distance of the point of libration to the center of gravity of two finite masses which are equal to 1 and μ , respectively. Since function Ω is a holomorphic function of ξ and η , at least in the direct proximity of the point of libration, it can be expanded into Taylor's series, converging absolutely for all values of ξ and η , the moduli of which do not exceed known limits. This series can be written in the form

$$\begin{aligned} \Omega = \frac{1}{2} p \xi^2 - \frac{1}{2} q \eta^2 - a \xi^3 + \frac{3}{2} a \xi \eta^2 + \beta \xi^4 - 3\beta \xi^2 \eta^2 \\ + \frac{3}{8} \beta \eta^4 - \gamma \xi^5 + 5\gamma \xi^3 \eta^2 - \frac{15}{8} \xi \eta^4 + \dots, \end{aligned} \quad (4)$$

where

$$\left. \begin{aligned} p &= 1 + \mu + 4f, \quad q = -1 - \mu + 2f, \\ 2f &= \frac{1}{\rho_{10}^3} + \frac{\mu}{\rho_{20}^3}, \quad a = \frac{1}{\rho_{10}^4} + \frac{\mu}{\rho_{20}^4}, \\ \beta &= \frac{1}{\rho_{10}^5} + \frac{\mu}{\rho_{20}^5}, \quad \gamma = \frac{1}{\rho_{10}^6} + \frac{\mu}{\rho_{20}^6}, \end{aligned} \right\} \quad (5)$$

whereby

$$\rho_{10} = \left| a - \frac{\mu}{1 + \mu} \right|, \quad \rho_{20} = \left| a + \frac{1}{1 + \mu} \right|. \quad (6)$$

If we retain only the terms of the second degree in our expansion of function Ω , Equation (1) will be transformed into linear equations with constant coefficients. The determining equation of this linear system, as is well known, has the form

$$x^4 - (p - q - 4n^2)x^2 - pq = 0. \quad (7)$$

This equation has the following roots:

$$\lambda_i, -\lambda_i, \lambda_1, -\lambda_1, \quad (i = \sqrt{-1}), \quad (8)$$

where

$$\left. \begin{aligned} \lambda_2 &= 1 + \mu - f + \sqrt{9f^2 - 4f(1 + \mu)}, \\ \lambda_1^2 &= -1 - \mu + f + \sqrt{9f^2 - 4f(1 + \mu)}. \end{aligned} \right\} \quad (9)$$

λ^2 always differs from zero and is positive. λ_1^2 becomes zero only with $\mu \neq 0$. Therefore, with all values of μ other than zero, Equation (7) possesses two purely imaginary and two real roots, one of which is positive and the other negative. The presence of one positive root of Equation (7) causes instability of collinear libration points in the Lyapunov sense, which is a well-known fact.

Let us now examine the matter of the existence of periodic motions near collinear points. This problem is solved without difficulty, with the aid of Lyapunov's theorem alone, proof for which can be found in his well known essay¹. This theorem can be formulated as follows: if the determining equation of a system of disturbed motion equations has two simple purely imaginary roots, whole multiples of which are not equation roots, and if the latter does not

possess zero roots, any time it is possible to find periodic series of a certain type which would formally satisfy the system of equations of disturbed motion, the latter, in fact, has a periodic solution presented by such series.

Thus, to find a periodic solution, it is necessary to set up series of which all terms are periodic functions. Therefore, in the general case, it is impossible to establish the existence of periodic solutions a priori, for it is virtually impossible to determine all terms of an infinite series.

Fortunately, there is at least one case whereby the existence of a periodic solution can be established a priori. This will be a case when equations of disturbed motion have a holomorphic integral, independent of time, of a definite structure. For example, if equations of disturbed motion have canonical form

$$\frac{dx_s}{dt} = -\frac{\partial H}{\partial y_s}, \quad \frac{dy_s}{dt} = \frac{\partial H}{\partial x_s}, \quad (s = 1, 2, \dots, n),$$

where H is a time-independent holomorphic function of variables x_s and y_s in which the terms of the lowest order generate a ground form, the equations have integral $H = \text{const.}$, and if the determining equation has a pair of simple purely imaginary roots λi and $-\lambda i$, not possessing roots of the type $m\lambda i$, where $m=0, \pm 1, \pm 2, \pm 3, \dots$, the disturbed motion equations will, without doubt, have a periodic solution represented by periodic series of known form.

Let us see what can be said of the existence of periodic motions near collinear libration points in the restricted problem of three bodies. As we have seen, the determining equation in this problem always has two simple purely imaginary roots and does not have (if $\mu \neq 0$) zero roots or roots of type $m\lambda i$, where m is a whole number. It is further known that system (1) has an integral independent of time. To become convinced that this integral has the required form, it is sufficient to reduce Equation (1) to canonical form, assuming

$$\xi_1 = \frac{d}{dt} - \eta \cdot \eta, \quad \eta_1 = \frac{d\eta}{dt} + \eta \cdot \xi.$$

Then Equation (1) will be replaced by the following system:

$$\left. \begin{aligned} \frac{d\xi}{dt} &= -\frac{\partial H}{\partial \xi_1}, & \frac{d\xi_1}{dt} &= \frac{\partial H}{\partial \xi}, \\ \frac{d\eta}{dt} &= -\frac{\partial H}{\partial \eta_1}, & \frac{d\eta_1}{dt} &= \frac{\partial H}{\partial \eta}, \end{aligned} \right\} \quad (10)$$

where

$$H = \Omega - \frac{n^2}{2} (\xi^2 + \eta^2) + n(\xi\eta_1 - \eta\xi_1) - \frac{1}{2} (\xi_1^2 + \eta_1^2). \quad (11)$$

Since H is independent of time, Equation (10) has a holomorphic integral in which the terms of the lowest order assume ground form H_2 , of type

$$H_2 = 2f\xi^2 - f\eta^2 + n(\xi\eta_1 - \eta\xi_1) - \frac{1}{2} (\xi_1^2 + \eta_1^2). \quad (12)$$

Therefore, system (10) indubitably has a periodic solution, and consequently, such a solution will exist for system (1).

Thus, we have established the existence of periodic motions near collinear points of libration, and it remains for us merely to find these motions. As Lyapunov² points out, for this we must proceed as follows. Assuming c to stand for arbitrary constant and T for series

$$T = \frac{2\pi}{\lambda} \left(1 + h_2 c^2 + h_3 c^3 + \dots \right) \quad (13)$$

with indefinite coefficients h_2, h_3, \dots , we shall insert in Equation (1) a new independent variable τ to replace t by substituting

$$\tau = \frac{2\pi(t - t_0)}{T}. \quad (14)$$

Constants h_2, h_3, \dots shall be selected in such a manner that transformed equation

$$\left. \begin{aligned} \frac{d^2 \xi}{d\tau^2} - 2n \frac{T}{2\pi} \cdot \frac{d\eta}{d\tau} &= \frac{T^2}{4\pi^2} \cdot \frac{\partial \Omega}{\partial \xi}, \\ \frac{d^2 \eta}{d\tau^2} + 2n \frac{T}{2\pi} \cdot \frac{d\xi}{d\tau} &= \frac{T^2}{4\pi^2} \cdot \frac{\partial \Omega}{\partial \eta}, \end{aligned} \right\} \quad (15)$$

will be satisfied by series

$$\left. \begin{aligned} \xi &= x_0 = cx_1 + c^2x_2 + c^3x_3 + \dots, \\ \eta &= y_0 = cy_1 + c^2y_2 + c^3y_3 + \dots, \end{aligned} \right\} \quad (16)$$

in which all coefficients, $x_1, y_1, x_2, y_2, \dots$, will be periodic functions of τ with common period 2π . Since the existence of a periodic solution in our problem is known a priori, such series will indeed be found and will be absolutely convergent at all values of τ and values of c satisfying condition

$$|c| \leq \bar{c}. \quad (17)$$

where \bar{c} is a certain positive value differing from zero. Series (13) will also converge absolutely with $|c| \leq \bar{c}$, and value T will be the period of the found motion (in respect to time). We have thus obtained a continuous series of periodic motions depending on two arbitrary constants, t_0 and c . Constant t_0 has no significant effect of the nature of the motion, since we can assume that the family of periodic motions depends solely on arbitrary constant c .

Determining in practice functions $x_1, y_1, x_2, y_2, \dots$, we shall obtain for them expressions of form

$$\left. \begin{aligned} x_1 &= A_1 [\cos \tau + \sin \tau], \\ y_1 &= B_1 [\cos \tau - \sin \tau], \\ x_2 &= A_2 \sin 2\tau + C_2, \\ y_2 &= B_2 \cos 2\tau \\ x_3 &= A_3 [\cos 3\tau - \sin 3\tau], \\ y_3 &= B_3 [\cos 3\tau + \sin 3\tau], \\ &\dots\dots\dots \end{aligned} \right\} \quad (18)$$

where $A_1, B_1, A_2, B_2, C_2, \dots$ are constants dependent only on μ . For example,

$$A_1 = \frac{2n}{\lambda}, \quad B_1 = 1 + \frac{p}{\lambda^2}.$$

We shall not write out expressions for the remaining coefficients, in view of their awkwardness.

Let us now turn to an investigation of the stability of the periodic motions found by us. In the absolute sense, this problem is solved immediately and in the negative. Indeed, let us examine any of the

found periodic motions. It corresponds to a certain value of constant c , and consequently, a certain period T . Since the family of periodic solutions is continuous in respect to c , there exists an innumerable quantity of other periodic motions orbitally as close as desired to that under examination. The periods of these orbitally close motions will differ from the period of the motion in question, but they will differ in a degree as small as desired. Let us now picture two particles located at the initial moment in two periodic orbits as close as desired, and possessing initial velocity coordinate and component values as close as desired. Let us say that T and T_1 are corresponding periods and let us assume, to be more definite, that $T < T_1$. It is obvious that after time interval T , the first particle will arrive at its initial state at that time when the second has not yet been able to reach that state. In other words, after time interval T , the second particle will lag somewhat behind the first and this gap will continue to grow with time. After a sufficiently large number of revolutions, our particles will be separated a distance comparable to the dimensions of their orbits and thus will cease being as close to one another as desired. This means that each periodic motion obtained by us is unstable in the Lyapunov sense.

But instability of this type is not particularly interesting to us. Indeed, the behavior of the other motions in our problem is still an unknown, that is, motions not belonging to our periodic family. It may occur that trajectories of all remaining motions, as close as desired to periodic according to initial data, will always be as close as desired. This will mean that the periodic motion is orbitally stable. On the other hand, it may occur that motions will be found in our problem as close to periodic as desired according to initial data, the trajectories of which will not remain as close as desired to periodic orbit, which will signify that this periodic motion is also orbitally unstable.

To elucidate this important matter of orbital stability of periodic motions near collinear points of libration we should examine Equation (15), in which τ is an independent variable. Indeed, in respect to τ , all periodic motions of the family possess common period 2π , as a consequence of which instability effect disappears (an effect occurring due to difference in periods of periodic orbits as close together as desired). We shall examine the problem of stability of any of solutions (16) of system (15). For this, we shall first insert new variables x and y in place of ξ and η , assuming

$$\xi = x + x_0, \quad \eta = y + y_0. \quad (19)$$

The transformed equations will have the form

$$\left. \begin{aligned} \frac{d^2 x}{d\tau^2} - 2n \frac{T}{2\pi} \cdot \frac{dy}{d\tau} &= \frac{T^2}{4\pi^2} \cdot \frac{\partial R}{\partial x}, \\ \frac{d^2 y}{d\tau^2} + 2n \frac{T}{2\pi} \cdot \frac{dx}{d\tau} &= \frac{T^2}{4\pi^2} \cdot \frac{\partial R}{\partial y}, \end{aligned} \right\} \quad (20)$$

where

$$R(x, y) = \Omega(x + x_0, y + y_0). \quad (21)$$

Since all periodic orbits lie in an area where function Ω is holomorphic, function $R(x, y)$ will also be a holomorphic function, at least for values of x and y , the moduli of which are sufficiently small. Therefore, function $R(x, y)$ can be represented in the form of a series with coefficients periodic in relation to τ . This series will be written as follows:

$$R = \frac{1}{2} p_{11}(\tau) x^2 + p_{12}(\tau) xy + \frac{1}{2} p_{22}(\tau) y^2 + \dots, \quad (22)$$

where the first coefficients will be determined by formulas

$$\left. \begin{aligned} p_{11}(\tau) &= p - 6\alpha x_0 + 12\beta x_0^2 - 6\beta_0^2 + \dots, \\ p_{12}(\tau) &= 3\alpha y_0 - 12\beta x_0 y_0 + \dots, \\ p_{22}(\tau) &= -q + 3\alpha x_0 - 6\beta x_0^2 + \frac{9}{2} \beta y_0^2 + \dots \end{aligned} \right\} \quad (23)$$

These coefficients are periodic functions of τ , with common period 2π and, in addition, are dependent on parameter c . Since these coefficients are values of partial derivatives of holomorphic function $\Omega(x + x_0, y + y_0)$ with $x=y=0$, series (23) will be absolutely convergent in the area where this function is holomorphic, and therefore coefficients $p_{11}(\tau), p_{12}(\tau), p_{22}(\tau), \dots$ can be represented in the form of series placed according to increasing degrees of parameter c and absolutely convergent at all values of τ and with $|c| \leq \bar{c}$. These series will have the form

$$\left. \begin{aligned} p_{11}(\tau) &= p + c p_{11}^{(1)}(\tau) + c^2 p_{11}^{(2)}(\tau) + \dots, \\ p_{12}(\tau) &= c p_{12}^{(1)}(\tau) + c^2 p_{12}^{(2)}(\tau) + \dots, \\ p_{22}(\tau) &= -q + c p_{22}^{(1)}(\tau) + c^2 p_{22}^{(2)}(\tau) + \dots, \end{aligned} \right\} \quad (24)$$

where all coefficients $p_{11}^{(1)}(\tau)$, $p_{12}^{(1)}(\tau)$, $p_{22}^{(1)}(\tau)$... are periodic functions of τ with common period 2π . Thus, our problem of periodic motion stability is reduced to a problem of stability of trivial equation

$$x = 0, \quad y = 0 \quad (ST)$$

of a system with periodic coefficients (20).

According to the general Lyapunov theory, we shall first examine the equations in variations corresponding to system (20). These equations in the variations will be written as follows:

$$\left. \begin{aligned} \frac{d^2 x}{d\tau^2} - 2n \frac{T}{2\pi} \frac{dy}{d\tau} &= \frac{T^2}{4\pi^2} [p_{11}(\tau)x + p_{12}(\tau)y], \\ \frac{d^2 y}{d\tau^2} + 2n \frac{T}{2\pi} \frac{dx}{d\tau} &= \frac{T^2}{4\pi^2} [p_{12}(\tau)x + p_{22}(\tau)y] \end{aligned} \right\} \quad (25)$$

and will represent a system of linear equations with periodic coefficients. The problem of stability (ST) of system (25) depends, as is known, on the nature of the roots of the characteristic equation of system (25). Since system (25) can be reduced to canonical form, its characteristic equation will have the form³

$$\rho^4 + A\rho^3 + B\rho^2 + A\rho + 1 = 0, \quad (26)$$

and coefficients A and B will be holomorphic functions of parameter c and can be represented in the form of series located by increasing degrees of c and absolutely convergent at $|c| \leq \tau^4$.

To determine the nature of the roots of Equation (24), we shall see what we shall have with $c = 0$. Assuming in Equation (25) $c = 0$, we shall obtain a system with constant coefficients

$$\left. \begin{aligned} \frac{d^2x}{d\tau^2} - \frac{2n}{\lambda} \frac{dy}{d\tau} - \frac{p}{\lambda^2} x &= 0, \\ \frac{d^2y}{d\tau^2} + \frac{2n}{\lambda} \frac{dx}{d\tau} + \frac{q}{\lambda^2} y &= 0, \end{aligned} \right\} \quad (27)$$

the determining equation of which will have the form

$$x^4 - \frac{1}{\lambda^2} (p - q - 4n^2) x^2 - \frac{pq}{\lambda^4} = 0. \quad (28)$$

Therefore, the roots of Equation (28) will be

$$i, \quad -i, \quad \frac{\lambda_2}{\lambda}, \quad -\frac{\lambda_1}{\lambda} \quad (i = \sqrt{-1}), \quad (29)$$

but system (27) can also be viewed as a system with periodic coefficients; therefore, for system (28), we can also build a characteristic equation of the same form as Equation (26). There is no need to set up the latter equation. The roots of the determining and characteristic equations of a system with constant coefficients are connected by formula⁵

$$\rho = e^{2\pi x}, \quad (30)$$

and since values of x are known, we immediately obtain corresponding values of ρ :

$$1, \quad 1, \quad e^{\frac{2\pi\lambda_1}{\lambda}}, \quad e^{-\frac{2\pi\lambda_1}{\lambda}}. \quad (31)$$

Hence, it is evident that with $c = 0$ Equation (26) possesses one root, the modulus of which is greater than one. Since A and B are holomorphic functions of c and since the above-mentioned root is simple, according to the theorem of the existence of implicit functions⁶, Equation (26)

with $c \neq 0$ will have a single root, tending to $e^{\frac{2\pi\lambda_1}{\lambda}}$, when c tends to zero. Therefore, with sufficiently small values of $|c|$ Equation (26) will obviously have a root the modulus of which is greater than one. According to Lyapunov's⁷ results, we conclude that (ST) of system (25) and (ST) of system (20) will be unstable.

This means that all periodic motions close to collinear points of libration in the restricted problem of three bodies also possess orbital instability, and the problem of stability of these periodic motions has thus been elucidated by us.

LITERATURE CITED

1. A. M. Lyapunov, GENERAL PROBLEMS OF STABILITY OF MOTION, ONTI, No. 42, 1935.
2. A. M. Lyapunov, GENERAL PROBLEMS OF STABILITY OF MOTION, ONTI, No. 36, 1935.
3. A. M. Lyapunov, GENERAL PROBLEMS OF STABILITY OF MOTION, ONTI, No. 51, 1935.
4. A. M. Lyapunov, GENERAL PROBLEMS OF STABILITY OF MOTION, ONTI, No. 48, 1935.
5. A. M. Lyapunov, GENERAL PROBLEMS OF STABILITY OF MOTION, ONTI, No. 46, 1935.
6. Gursa, COURSE ON MATHEMATICAL ANALYSIS (Kurs Matematicheskogo Analiza), Vol. 1.
7. A. M. Lyapunov, GENERAL PROBLEMS OF STABILITY OF MOTION, ONTI, No. 55, 1935.

DISTRIBUTION

	No. of Copies		No. of Copies
<u>EXTERNAL</u>		U. S. Atomic Energy Commission	1
Air University Library	1	ATTN: Reports Library, Room G-017	
ATTN: AUL3T		Washington, D. C. 20545	
Maxwell Air Force Base, Alabama 36112		U. S. Naval Research Laboratory	1
U. S. Army Electronics Proving Ground	1	ATTN: Code 2027	
ATTN: Technical Library		Washington, D. C. 20390	
Fort Huachuca, Arizona 85613		Weapons Systems Evaluation Group	1
U. S. Naval Ordnance Test Station	1	Washington, D. C. 20305	
ATTN: Technical Library, Code 753		John F. Kennedy Space Center, NASA	2
China Lake, California 93555		ATTN: KSC Library, Documents Section	
U. S. Naval Ordnance Laboratory	1	Kennedy Space Center, Florida 32899	
ATTN: Library		APGC (PGBPS-12)	1
Corona, California 91720		Eglin Air Force Base, Florida 32542	
Lawrence Radiation Laboratory	1	U. S. Army CDC Infantry Agency	1
ATTN: Technical Information Division		Fort Benning, Georgia 31905	
P. O. Box 808		Argonne National Laboratory	1
Livermore, California 94550		ATTN: Report Section	
Sandia Corporation	1	9700 South Cass Avenue	
ATTN: Technical Library		Argonne, Illinois 60440	
P. O. Box 969		U. S. Army Weapons Command	1
Livermore, California 94551		ATTN: AMSWE-RDR	
U. S. Naval Postgraduate School	1	Rock Island, Illinois 61201	
ATTN: Library		Rock Island Arsenal	1
Monterey, California 93940		ATTN: SWERI-RDI	
Electronic Warfare Laboratory, USAECOM	1	Rock Island, Illinois 61201	
Post Office Box 205		U. S. Army Cmd. & General Staff College	1
Mountain View, California 94042		ATTN: Acquisitions, Library Division	
Jet Propulsion Laboratory	2	Fort Leavenworth, Kansas 66027	
ATTN: Library (TDS)		Combined Arms Group, USACDC	1
4800 Oak Grove Drive		ATTN: Op. Res., P and P Div.	
Pasadena, California 91103		Fort Leavenworth, Kansas 66027	
U. S. Naval Missile Center	1	U. S. Army CDC Armor Agency	1
ATTN: Technical Library, Code N3022		Fort Knox, Kentucky 40121	
Point Mugu, California 93041		Michoud Assembly Facility, NASA	1
U. S. Army Air Defense Command	1	ATTN: Library, I-MICH-OSD	
ATTN: ADSX		P. O. Box 29300	
Ent Air Force Base, Colorado 80912		New Orleans, Louisiana 70129	
Central Intelligence Agency	4	Aberdeen Proving Ground	1
ATTN: OCR/DD-Standard Distribution		ATTN: Technical Library, Bldg. 313	
Washington, D. C. 20505		Aberdeen Proving Ground, Maryland 21005	
Harry Diamond Laboratories	1	NASA Sci. & Tech. Information Facility	5
ATTN: Library		ATTN: Acquisitions Branch (S-AK/DL)	
Washington, D. C. 20438		P. O. Box 33	
Scientific & Tech. Information Div., NASA	1	College Park, Maryland 20740	
ATTN: ATS		U. S. Army Edgewood Arsenal	1
Washington, D. C. 20546		ATTN: Librarian, Tech. Info. Div.	
		Edgewood Arsenal, Maryland 21010	

	No. of Copies		No. of Copies
National Security Agency ATTN: C3/TDL Fort Meade, Maryland 20755	1	Brookhaven National Laboratory Technical Information Division ATTN: Classified Documents Group Upton, Long Island, New York 11973	1
Goddard Space Flight Center, NASA ATTN: Library, Documents Section Greenbelt, Maryland 20771	1	Watervliet Arsenal ATTN: SWEV-RD Watervliet, New York 12189	1
U. S. Naval Propellant Plant ATTN: Technical Library Indian Head, Maryland 20640	1	U. S. Army Research Office (ARO-D) ATTN: CRD-AA-IP Box CM, Duke Station Durham, North Carolina 27706	1
U. S. Naval Ordnance Laboratory ATTN: Librarian, Eva Liberman Silver Spring, Maryland 20910	1	Lewis Research Center, NASA ATTN: Library 21000 Brookpark Road Cleveland, Ohio 44135	1
Air Force Cambridge Research Labs. L. G. Hanscom Field ATTN: CRMCLR/Stop 29 Bedford, Massachusetts 01730	1	Systems Engineering Group (RTD) ATTN: SEPIR Wright-Patterson Air Force Base, Ohio 45433	1
Springfield Armory ATTN: SWESP-RE Springfield, Massachusetts 01101	1	U. S. Army Artillery & Missile School ATTN: Guided Missile Department Fort Sill, Oklahoma 73503	1
U. S. Army Materials Research Agency ATTN: AMXMR-ATL Watertown, Massachusetts 02172	1	U. S. Army CDC Artillery Agency ATTN: Library Fort Sill, Oklahoma 73504	1
Strategic Air Command (OAI) Offutt Air Force Base, Nebraska 68113	1	U. S. Army War College ATTN: Library Carlisle Barracks, Pennsylvania 17013	1
Picatinny Arsenal, USAMUCOM ATTN: SMUPA-VA6 Dover, New Jersey 07801	1	U. S. Naval Air Development Center ATTN: Technical Library Johnsville, Warminster, Pennsylvania 18974	1
U. S. Army Electronics Command ATTN: AMSEL-CB Fort Monmouth, New Jersey 07703	1	Frankford Arsenal ATTN: C-2500-Library Philadelphia, Pennsylvania 19137	1
Sandia Corporation ATTN: Technical Library P. O. Box 5800 Albuquerque, New Mexico 87115	1	Div. of Technical Information Ext., USAEC P. O. Box 62 Oak Ridge, Tennessee 37830	1
ORA(RRRT) Holloman Air Force Base, New Mexico 88330	1	Oak Ridge National Laboratory ATTN: Central Files P. O. Box X Oak Ridge, Tennessee 37830	1
Los Alamos Scientific Laboratory ATTN: Report Library P. O. Box 1663 Los Alamos, New Mexico 87544	1	Air Defense Agency, USACDC ATTN: Library Fort Bliss, Texas 79916	1
White Sands Missile Range ATTN: Technical Library White Sands, New Mexico 88002	1	U. S. Army Air Defense School ATTN: AKBAAS-DR-R Fort Bliss, Texas 79906	1
Rome Air Development Center (EMLAL-1) ATTN: Documents Library Griffiss Air Force Base, New York 13440	1		

	No. of Copies		No. of Copies
U. S. Army CDC Nuclear Group Fort Bliss, Texas 79916	1	<u>INTERNAL</u>	
Manned Spacecraft Center, NASA ATTN: Technical Library, Code BM6 Houston, Texas 77058	1	Headquarters U. S. Army Missile Command Redstone Arsenal, Alabama ATTN: AMSMI-D	1
Defense Documentation Center Cameron Station Alexandria, Virginia 22314	20	AMSMI-XE, Mr. Lowers	1
		AMSMI-XS, Dr. Carter	1
		AMSMI-Y	1
		AMSMI-R, Mr. McDaniel	1
		AMSMI-RAP	1
U. S. Army Research Office ATTN: STINFO Division 3045 Columbia Pike Arlington, Virginia 22204	1	AMSMI-RBLD	10
		USACDC-LnO	1
		AMSMI-RB, Mr. Croxton	1
		AMSMI-RBT	8
U. S. Naval Weapons Laboratory ATTN: Technical Library Dahlgren, Virginia 22448	1	National Aeronautics & Space Administration Marshall Space Flight Center Huntsville, Alabama	
U. S. Army Engineer Res. & Dev. Labs. ATTN: Scientific & Technical Info. Br. Fort Belvoir, Virginia 22060	2	ATTN: MS-T, Mr. Wiggins	5
		R-AERO-X, Mr. Wheeler	1
Langley Research Center, NASA ATTN: Library, MS-185 Hampton, Virginia 23365	1		
Research Analysis Corporation ATTN: Library McLean, Virginia 22101	1		
U. S. Army Tank Automotive Center ATTN: SMOTA-RTS.1 Warren, Michigan 48090	1		
Hughes Aircraft Company Electronic Properties Information Center Florence Ave. & Teale St. Culver City, California 90230	1		
Atomics International, Div. of NAA Liquid Metals Information Center P. O. Box 309 Canoga Park, California 91305	1		
Foreign Technology Division ATTN: Library Wright-Patterson Air Force Base, Ohio 45400	1		
Clearinghouse for Federal Scientific and Technical Information U. S. Department of Commerce Springfield, Virginia 22151	1		
Foreign Science & Technology Center, USAMC ATTN: Mr. Shapiro Washington, D. C. 20315	3		

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R&D		
<i>(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)</i>		
1. ORIGINATING ACTIVITY <i>(Corporate author)</i> Redstone Scientific Information Center Research and Development Directorate U. S. Army Missile Command Redstone Arsenal, Alabama 35809		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP N7A
3. REPORT TITLE REPORT 4: INSTABILITY OF PERIODIC MOTIONS NEAR COLLINEAR POINTS OF LIBRATION IN THE RESTRICTED PROBLEM OF THREE BODIES Astronomicheskii Zhurnal, 15, No. 3, 209-216 (1938)		
4. DESCRIPTIVE NOTES <i>(Type of report and inclusive dates)</i> Translated from the Russian		
5. AUTHOR(S) <i>(Last name, first name, initial)</i> Duboshin, G. N.		
6. REPORT DATE 13 February 1967	7a. TOTAL NO. OF PAGES 14	7b. NO. OF REFS 7
8a. CONTRACT OR GRANT NO. N/A	9a. ORIGINATOR'S REPORT NUMBER(S) RSIC -647	
b. PROJECT NO. N/A		
c.	9b. OTHER REPORT NO(S) <i>(Any other numbers that may be assigned this report)</i>	
d.	AD _____	
10. AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited.		
11. SUPPLEMENTARY NOTES None	12. SPONSORING MILITARY ACTIVITY Same as No. 1	
13. ABSTRACT Examined is the problem of periodic motions near collinear points of libration in the plane restricted problem of three bodies. With the aid of the Lyapunov methods, it is demonstrated that around each collinear point of libration a continuous family of periodic motions exists, dependent on one arbitrary constant. Examined is the stability of the indicated periodic motions; it is shown that they are unstable. Also established is the orbital instability of the periodic motions under study.		

DD FORM 1 JAN 64 1473

UNCLASSIFIED

Security Classification

15

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Periodic motions Collinear points of libration Taylor's series Holomorphic integral Lyapunov theory						

INSTRUCTIONS

1. ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. REPORT DATE: Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. NUMBER OF REFERENCES: Enter the total number of references cited in the report.

8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. AVAILABILITY/LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.

12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.